

Renormalization group approach to satisfiability

S.N. Coppersmith

*Department of Physics, University of Wisconsin,
1150 University Avenue, Madison, WI 53706 USA*

(Dated: February 2, 2008)

Satisfiability is a classic problem in computational complexity theory, in which one wishes to determine whether an assignment of values to a collection of Boolean variables exists in which all of a collection of clauses composed of logical OR's of these variables is true. Here, a renormalization group transformation is constructed and used to relate the properties of satisfiability problems with different numbers of variables in each clause. The transformation yields new insight into phase transitions delineating “hard” and “easy” satisfiability problems.

PACS numbers: 05.10.Cc, 89.20.Ff, 75.10.Nr

Computational complexity theory addresses the question of how fast the resources required to solve a given problem grow with the size of the input needed to specify the problem.[1] P is the class of problems that can be solved in polynomial time, which means a time that grows as a polynomial of the size of the problem specification, while NP is the class of problems for which a solution can be verified in polynomial time. Whether or not P is distinct from NP has been a central unanswered question in computational complexity theory for decades.[2]

Satisfiability (SAT) is a classic problem in computational complexity. An often-studied type of SAT is K -SAT, in which one attempts to find assignment of N variables such that the conjunction (AND) of M constraints, or clauses, each of which is the disjunction (OR) of K literals, each literal being either a negated or un-negated variable, is true. (This way of writing the problem, as a conjunction of clauses that are disjunctions, is called conjunctive normal form.) For example, the 3-SAT instance with the 4 variables x_1, x_2, x_3 , and x_4 and the four clauses

$$\begin{aligned} & (x_1 = 1 \text{ OR } x_2 = 0 \text{ OR } x_4 = 1) \\ \text{AND} & \quad (x_1 = 0 \text{ OR } x_3 = 1 \text{ OR } x_4 = 0) \\ \text{AND} & \quad (x_2 = 1 \text{ OR } x_3 = 1 \text{ OR } x_4 = 1) \\ \text{AND} & \quad (x_1 = 0 \text{ OR } x_3 = 0 \text{ OR } x_4 = 1), \end{aligned} \quad (1)$$

is satisfiable because it is true for the assignments $x_1 = 1, x_2 = 1, x_3 = 1, x_4 = 1$. Below, we will write satisfiability problems in conjunctive normal form using the notation of [3], where the ANDs and ORs are implied and the literals have positive or negative signs depending on whether or not they are negated. For example, the expression of Eq. (1) is written

$$(1 \text{ } -2 \text{ } 4), (-1 \text{ } 3 \text{ } -4), (2 \text{ } 3 \text{ } 4), (-1 \text{ } -3 \text{ } 4).$$

2-SAT can be solved in polynomial time [1], while K -SAT with $K \geq 3$ is known to be NP-complete [4]: if a polynomial algorithm for 3-SAT exists, then P is equal to NP. The complexity of SAT is intimately related to the presence of phase transitions [5, 6, 7, 8, 9, 10, 11]. For random problems with N variables, M clauses, and K literals per clause, as M is increased there is a phase transition from a satisfiable phase, in which almost all random instances are satisfiable, to an unsatisfiable phase, in which almost all random instances are unsatisfiable. The most difficult instances are near this SAT-unSAT transition. It has also been shown that there is a transition as the parameter K is changed between 2 and 3, at $K_c \sim 2.4$, at which the nature of the SAT-unSAT transition changes [5].

Here, we investigate the relationship between satisfiability problems with different values of K by constructing a renormalization group transformation, similar to those used for phase transition problems [12, 13, 14, 15], that reduces the number of degrees of freedom, while possibly increasing the number and range of interactions [16] (which in this context is the number of literals per clause). To do this, we note that the expression

$$\begin{aligned} & ((\mathcal{A}_1 x), (\mathcal{A}_2 x), \dots (\mathcal{A}_P x), \\ & (\mathcal{B}_1 - x), (\mathcal{B}_2 - x)) \dots, (\mathcal{B}_Q - x)) \end{aligned}$$

is satisfiable if and only if

$$\begin{aligned} & ((\mathcal{A}_1 \mathcal{B}_1), (\mathcal{A}_1 \mathcal{B}_2), \dots (\mathcal{A}_1 \mathcal{B}_Q), \\ & (\mathcal{A}_2 \mathcal{B}_1), (\mathcal{A}_2 \mathcal{B}_2), \dots (\mathcal{A}_2 \mathcal{B}_Q), \\ & \dots, \\ & (\mathcal{A}_P \mathcal{B}_1), (\mathcal{A}_P \mathcal{B}_2), \dots, (\mathcal{A}_P \mathcal{B}_Q)) \end{aligned}$$

is satisfiable. Here, the $\mathcal{A}_i$'s and $\mathcal{B}_i$'s are arbitrary clauses and x is a variable. (The easiest way to see the equivalence is to note that both expressions are satisfiable if and only if $(\mathcal{A}_1 \text{ AND } \mathcal{A}_2 \text{ AND } \dots \mathcal{A}_P) \text{ OR } (\mathcal{B}_1 \text{ AND } \mathcal{B}_2 \text{ AND } \dots \mathcal{B}_Q)$ is.) The first step of the renormalization procedure is to use this identity to eliminate a given variable. In this step, P clauses in which a given variable comes in un-negated and Q clauses in which the same variable comes in negated are eliminated and replaced with PQ "resolution" [17] clauses. Thus, eliminating a "frustrated" [18] variable (one that enters into different clauses negated and un-negated) increases the number of clauses if $PQ - (P+Q) > 0$. The resolution of two clauses of length K_i and K_j has length $K_i + K_j - 2$. Note that resolving two 2-clauses yields a 2-clause, resolving a 2-clause with a clause of length $K \geq 3$ yields a clause length K , and resolving two clauses of with lengths $K_1 \geq 3$ and $K_2 \geq 3$ yields a clause with length greater than both K_1 and K_2 .

One then simplifies the resulting satisfiability expression by noting that

1. Duplicate clauses are redundant,
2. Duplicate literals in a given clause are redundant,
3. If a variable enters into one clause both negated and un-negated, then the clause must be true and can be removed,
4. If a clause has one literal, then the value of the corresponding variable is determined, and
5. If a subset of the literals in a clause comprise a different clause, then the clause with more literals is redundant.

This last point means, for example, that if an expression contains both $(1 \ 3 \text{ -}4 \ 5)$ and $(1 \ 3)$, then $(1 \ 3 \text{ -}4 \ 5)$ can be removed, because it is satisfied automatically if $(1 \ 3)$ is satisfied.

This procedure is known in computer science as "the Davis-Putnam procedure of 1960 [19] with subsumption [20]," and was originally proposed as a method for solving satisfiability instances. It does not perform well in practice [21], and has been proven to require exponential time on some instances [22, 23]. However, here the aim is not to solve a given instance, but rather to investigate the "flow" of the problem itself as variables are eliminated [12, 24]. In particular, this renormalization group (RG) transformation provides a natural

framework for understanding a phase transitions between "easy" and "hard" satisfiability problems identified in [5].

We present evidence that the change in the nature of the SAT-unSAT phase transition critical value $K_c \sim 2.4$ [5] is intimately related to whether or not the number of clauses proliferates exponentially upon repeated application of the renormalization group (RG) transformation. Note that when $K = 2$ the clause length decreases upon renormalization, since the resolution of two 2-clauses is a 2-clause, so no clause gets longer, and some of the resulting clauses have a duplicate literal and so get shorter. Having a large number of 2-clauses limits the growth in the number of long clauses because of subsumption, so there is a qualitative difference in the behavior depending on whether the ratio of the number of 2-clauses to the number of variables grows or shrinks upon renormalization.

We show numerical data for an RG implementation in which successive variables are chosen randomly and eliminated if they occur in a clause of minimum length. This procedure is used because it focuses on short clauses, which are much more restrictive than long clauses. Figure 1 shows α_K , the ratio of M_K , the number of clauses of length K to N , the number of variables remaining in the problem, as a function of K , as the RG proceeds. The average and standard deviation of numerical data from 5 realizations at the SAT-unSAT transition with $p = 0.2$ and $p = 0.6$ are shown (using parameter values for the transition locations from [5]). Large numbers of long clauses are generated when $p = 0.6 > p_c$ and not when $p = 0.2 < p_c$.

It has been proven that the SAT-unSAT transition for 2-SAT occurs at $\alpha = 1$ [25], and for $2 \leq K < 2.4$, the SAT-unSAT transition is believed to occur when $\alpha_2 = 1$ [5]. Figure 2 (left) shows that when $K = 2.2$, α_2 increases upon renormalization when $\alpha_2 > 1$ and decreases upon renormalization when $\alpha_2 < 1$. Because adding additional three-clauses does not affect the behavior of the two-clauses, and because two-clauses are much more restrictive than longer clauses, the two-clauses dominate the problem whenever $\alpha_2 > 1$. One would expect that 3-clauses by themselves would proliferate when $N_{initial}$, the initial number of variables, was equal to $3M_3/2$. (This estimate, the analog of the result for two-clauses, follows from setting the number of literals in all 3-clauses to twice the number of variables, which means that on average each variable enters




FIG. 1: Plot of α_K , the ratio of M_K , the number of clauses of length K , to N , the number of variables, at different stages of the renormalization process. The points plotted are the mean and standard deviation of the results from five independent system realizations. The parameters are chosen to be at the SAT-unSAT transition with $p = 0.2 < p_c$ (left panel) and $p = 0.6 > p_c$ (right panel). When $p > p_c$ the clause length increases markedly and the number of clauses grows enormously.

into one 3-clause negated and one 3-clause un-negated. Moreover, eliminating one variable on average yields two less literals and one less variable, so that, ignoring fluctuations, the relationship remains true.) However, because the 2-clauses prevent the 3-clauses from proliferating, adding 3-clauses to the 2-clauses changes the nature of the SAT-unSAT transition only when enough 3-clauses have been added so that the SAT-unSAT transition occurs with $\alpha_2 < 1$. In this regime, under renormalization the 2-clauses disappear and so the clauses all become longer. Since very long clauses are ORs of many literals and hence easy to satisfy, the number of clauses must go up sufficiently fast for the problem to be difficult to solve—at large K , the SAT-unSAT transition occurs when the ratio of the number of clauses to the number of variables is $\propto 2^K$ [11]. When $K > K_c$, near the SAT-unSAT transition we expect the number of clauses to grow geometrically with iteration number, and the numerical data are consistent with the maximum number of clauses obtained during the renormalization process increasing exponentially with the initial number of variables, with an exponent that increases monotonically with $(M/N)_{\text{initial}}$, the initial ratio of the number of clauses to the number of variables. Figure 2 (right) shows a phase boundary lines for the SAT-




FIG. 2: Left: plot of α_2 , the ratio of the number of 2-clauses to the number of variables, versus N , the number of undecimated variables, for instances with $p = 0.2$ and different initial values of α . The numerical data are averages and standard deviations of five realizations of systems of size 500 when the initial $\alpha = 1$, size 400 when the initial $\alpha = \alpha_c = 1.2$, and size 300 when the initial $\alpha = 1.67$. The numerical data are consistent with the hypothesis that when $K < K_c$ the SAT-unSAT transition occurs when α_2 neither decreases nor increases upon renormalization. Right: Schematic phase diagram showing the SAT-unSAT transition (using data of Ref. [5]), the region in which the number of 2-clauses increases upon renormalization (the red hatched region in the left of the figure) and an estimate of the region in which the number of clauses with $K \geq 3$ increases upon renormalization (the blue hatched region in the right of the figure). The SAT-unSAT transition line crosses into the region in which long clauses proliferate exponentially at the intersection of the three lines. The estimate for 3-clause proliferation given in the text yields an intersection at $(K = 2.4, \alpha = 1.25)$.

unSAT transition, for the onset of increase in the number of 2-clauses, and our estimate for the onset of proliferation of clauses of length greater than or equal to three.

When $K > K_c$, the SAT-unSAT transition [5, 6, 7, 8, 9, 10, 26] occurs in a regime in which the renormalization transformation causes both the typical clause length and the total number of clauses to grow. We conjecture that the SAT-UNSAT transition occurs at the value of M/N at which the rate of exponential growth is a critical value. A “replica-symmetry breaking” transition [7, 9, 10, 26] at a somewhat smaller value of α can be interpreted in terms of propagation of constraints on eliminated literals, as will be discussed elsewhere. [27] Because of the exponential clause pro-

liferation, numerical investigation of these transitions using this renormalization group is limited to small sizes. However, the renormalization group may still be useful for investigating these transitions using analytic techniques appropriate for large K [10, 11] for any $K > K_c$, though it will be necessary to understand how to account for possible RG-induced correlations between clauses.

When $K > K_c$, the number of clauses continues to increase under renormalization until it is no longer unlikely that a given compound clause contains a repeated variable (in addition to the decimated one), which we expect to occur when the renormalized clause length is of order $\sqrt{N}$ [28]. Because it appears that there is no impediment to the growth in the effective value of K until it is of order $\sqrt{N}$, where the problem specification itself is exponentially large in N , it appears that the renormalization group procedure can transform the problem out of NP and even PSPACE altogether. This property may indicate that whether or not a given computational problem has a solution that can be verified using polynomially-bounded resources has no fundamental effect on the difficulty of solving the problem.

In summary, a transformation inspired by the renormalization group is constructed and used to relate the behavior of satisfiability problems with different values of K , the number of literals per clause. The transformation provides useful insight into previously identified phase transitions of satisfiability problems and may yield new insight into the question of whether or not P is equal to NP.

The author gratefully acknowledges financial support from grants nsf-dmr 0209630 and nsf-emt 0523680, and the hospitality of the Aspen Center for Physics, where some of this work was done.

[1] C. Papadimitriou, *Computational Complexity* (Addison-Wesley, 1994).

[2] Clay Mathematics Institute Millenium Problems; <http://www.claymath.org/millennium/>.

- [3] SATLIB - The Satisfiability Library; <http://www.satlib.org>.
- [4] S. Cook, in *Proceedings of the third annual ACM symposium on Theory of computing* (ACM, New York, 1971), pp. 151–158.
- [5] R. Monasson, R. Zecchina, S. Kirkpatrick, B. Selman, and L. Troyansky, *Nature* **400**, 133 (1999).
- [6] S. Kirkpatrick and B. Selman, *Science* **264**, 1297 (1994).
- [7] M. Mézard, G. Parisi, and R. Zecchina, *Science* **297**, 812 (2002).
- [8] G. Biroli, R. Monasson, and M. Weigt, *Eur. Phys. J. B* **14**, 551 (2000).
- [9] M. Mézard and R. Zecchina, *Phys. Rev. E* **66**, 056126 (2002).
- [10] T. Mora, M. Mézard, and R. Zecchina (2005), preprint cond-mat/0506053.
- [11] D. Achlioptas, A. Naor, and Y. Peres, *Nature* **435**, 759 (2005).
- [12] K. Wilson, *Phys. Rev. B* **4**, 3174 (1970).
- [13] K. Wilson, *Scientific American* **241**, 158 (1979).
- [14] H. Maris and L. Kadanoff, *Am. J. Phys.* **46**, 652 (1978).
- [15] S. White, *Phys. Rev. Lett.* **69**, 2863 (1992).
- [16] T. Niemeijer and J. V. Leeuwen, in *Phase transitions and critical phenomena*, vol. 6 (Academic, New York, 1976).
- [17] J. Robinson, *J. Assoc. Comput. Machin.* **12**, 23 (1965).
- [18] G. Toulouse, *Phys. Rep.* **49**, 267 (1979).
- [19] M. Davis and H. Putnam, *J. of the ACM* **7**, 201 (1960).
- [20] S. Cook and R. Reckhow, *J. Symbolic Logic* **44**, 36 (1979).
- [21] L. Zhang and S. Malik, in *Lecture Notes in Computer Science*, Vol. 2404 (Springer-Verlag, London, 2002), pp. 17–36.
- [22] G. Tseitin, in *Studies in Constructive Mathematics and Mathematical Logic, Part II*, edited by A. Slisenko (Consultants Bureau, 1968), pp. 115–125.
- [23] A. Haken, *Theor. Comp. Sci.* **39**, 297 (1985).
- [24] L. Kadanoff, *Physics* **2**, 263 (1966).
- [25] B. Bollobas, C. Borgs, J. Chayes, J. Kim, and D. Wilson, *Random Structures and Algorithms* **18**, 201 (2001).
- [26] S. Mertens, M. Mézard, and R. Zecchina (2005), <http://arxiv.org/abs/cs.CC/0309020>.
- [27] S. Coppersmith, unpublished.
- [28] P. Diaconis and F. Mosteller, *J. Amer. Statist. Assoc.* **84**, 853 (1989).